

UNCLASSIFIED

Defense Technical Information Center
Compilation Part Notice

ADP014937

TITLE: Plasma Phase Transition in Dense Hydrogen and Electron-Hole Plasmas

DISTRIBUTION: Approved for public release, distribution unlimited

This paper is part of the following report:

TITLE: International Conference on Phenomena in Ionized Gases [26th]
Held in Greifswald, Germany on 15-20 July 2003. Proceedings, Volume 4

To order the complete compilation report, use: ADA421147

The component part is provided here to allow users access to individually authored sections of proceedings, annals, symposia, etc. However, the component should be considered within the context of the overall compilation report and not as a stand-alone technical report.

The following component part numbers comprise the compilation report:

ADP014936 thru ADP015049

UNCLASSIFIED

Plasma Phase Transition in Dense Hydrogen and Electron-Hole Plasmas

V.S. Filinov¹, M. Bonitz², P. Levashov¹, V.E. Fortov¹, W. Ebeling³ and M. Schlanges⁴

¹ Institute for High Energy Density, Russian Academy of Sciences, Izhoroskay 13/19, Moscow 127412, Russia

² Fachbereich Physik, Universität Rostock, D-18051 Rostock, Germany

³ Institut für Physik, Humboldt-Universität Berlin, Invalidenstrasse 110, D-10115 Berlin, Germany

⁴ Fachbereich Physik, Universität Greifswald, D-17489 Greifswald, Germany

Plasma phase transitions (PPT) in dense hydrogen and electron-hole plasmas are investigated by direct path integral Monte Carlo methods (DPIMC). The phase boundary of the electron-hole liquid in Germanium is calculated and is found to agree reasonably well with the known experimental results. Analogous results are found for high-density hydrogen. For a temperature of $T = 10,000\text{K}$ it is shown that the internal energy is lowered due to droplet formation for densities between 10^{23}cm^{-3} and 10^{24}cm^{-3} .

1. Path integral Monte Carlo simulations

All thermodynamic properties of a two-component plasma are defined by the partition function Z which, for the case of N_e electrons and N_p protons, is given by $Z(N_e, N_p, V, \beta) = \frac{Q(N_e, N_p, \beta)}{N_e! N_p!}$, with $Q(N_e, N_p, \beta) = \sum_{\sigma} \int d\mathbf{q} d\mathbf{r} \rho(\mathbf{q}, \mathbf{r}, \sigma; \beta)$, where $\beta = 1/k_B T$. The exact density matrix is, for a quantum system, in general, not known but can be constructed using a path integral representation $[1, 2] \int_V dR^{(0)} \sum_{\sigma} \rho(R^{(0)}, \sigma; \beta) = \int_V dR^{(0)} \dots dR^{(n)} \rho^{(1)} \cdot \rho^{(2)} \dots \rho^{(n)} \times \sum_{\sigma} \sum_P (\pm 1)^{\kappa_P} S(\sigma, \hat{P}\sigma') \hat{P} \rho^{(n+1)}$, where $\rho^{(i)} \equiv \rho(R^{(i-1)}, R^{(i)}; \Delta\beta) \equiv \langle R^{(i-1)} | e^{-\Delta\beta \hat{H}} | R^{(i)} \rangle$, whereas $\Delta\beta \equiv \beta/(n+1)$ and $\Delta\lambda_a^2 = 2\pi\hbar^2 \Delta\beta/m_a$, $a = p, e$. \hat{H} is the Hamilton operator, $\hat{H} = \hat{K} + \hat{U}_c$, containing kinetic and potential energy contributions, \hat{K} and \hat{U}_c , respectively, with $\hat{U}_c = \hat{U}_p^e + \hat{U}_e^e + \hat{U}_e^p$ being the sum of the Coulomb potentials between protons (p), electrons (e) and electrons and protons (ep). Further, σ comprises all particle spins, and

the particle coordinates are denoted by $R^{(i)} = (q^{(i)}, r^{(i)}) \equiv (R_p^{(i)}, R_e^{(i)})$, for $i = 1, \dots, n+1$, $R^{(0)} \equiv (q, r) \equiv (R_p^{(0)}, R_e^{(0)})$, and $R^{(n+1)} \equiv R^{(0)}$ and $\sigma' = \sigma$. This means, the particles are represented by fermionic loops with the coordinates (beads) $[R] \equiv [R^{(0)}; R^{(1)}; \dots; R^{(n)}; R^{(n+1)}]$, where q and r denote the electron and proton coordinates, respectively. The spin gives rise to the spin part of the density matrix S , whereas exchange effects are accounted for by the permutation operator \hat{P} , which acts on the electron coordinates and spin projections, and the sum over the permutations with parity κ_P . To compute thermodynamic functions, the logarithm of the partition function has to be differentiated with respect to thermodynamic variables, so for internal energy E we have $\beta E = -\beta \ln Q / \partial \beta$

2. Numerical Results

Since the PPT in dense hydrogen is still hypothetical and has not been observed experimentally, it is reasonable to look for other systems where similar conditions exist. A suitable example is electron-hole plasma in low-temperature semiconductors, for which droplet formation is well established and observed ex-

perimentally three decades ago [3]. We, therefore, performed DPIMC simulations for electron hole plasmas. Below the critical temperature the simulations exhibit anomalously large fluctuations and an unstable behavior of the pressure. The e-h-plasma is found to phase separate and form large droplets. The phase boundary of the electron-hole liquid (e-h-droplets) in Germanium obtained by our DPIMC method is presented in Fig. 1 together with the experimental data. We observe good agreement. Deviations may be connected with complex band structure of Germanium approximated in our simulations by a two-band parabolic mass model.




Figure 1: Phase boundary of the electron-hole liquid in bulk Germanium. Experiment - [3]. Temperature is presented in units of the exciton binding energy.

Fig.2 presents results of our calculations for pressure and energy of hydrogen at $T = 10,000K$, which is well below the critical point of the PPT predicted by chemical models (around $T = 15,000K$). In our calculations between $10^{22}cm^{-3}$ and $10^{24}cm^{-3}$ calculated pressure becomes negative. We find also that in this region plasma energy is systematically lower than the RPIMC results [4]. If average distance be-




Figure 2: Pressure and energy of hydrogen for $T = 10,000K$, DFT - [5], RPIMC - [4].

tween plasma particles is of the order of the size of a hydrogen molecule the homogeneous plasma state becomes unstable, and many-particle clusters appear. Results of one independent well tested method based on density functional theory (DFT) are presented on Fig. 2, where PPT was obtained at smaller density [5].

References

- [1] V.S.Filinov, M.Bonitz, W.Ebeling, V.Fortov, Plasma Phys. Contr. Fusion **43** (2001) 743.
- [2] V.Filinov, J.Phys.A **34** (2001) 1665.
- [3] G.Thomas, T.Rice, J.Hensel, Phys.Rev.Let., **33** (1974) 219.
- [4] B.Militzer, D.Ceperley, Phys. Rev. Lett. **85** (2000) 1890.
- [5] H.Xu, J.Hansen, Phys. Rev.E **57** (1998) 211.